

Kaczmarek K., Brylewski P., Szymankiewicz-Szukała A., Wincek A. *Current neurophysiological methods for the assessment of motor function and therapy in patients treated for ischemic stroke. Issue Rehabil. Orthop. Neurophysiol. Sport Promot. 2019; 26: xxx-xxx. DOI: 10.19271/IRONS-000087-2019-26*

AKTUALNE METODY NEUROFIZJOLOGICZNE OCENY FUNKCJI RUCHOWEJ I TERAPII U CHORYCH LECZONYCH Z POWODU UDARU NIEDOKRWIENNEGO

Katarzyna Kaczmarek¹, Piotr Brylewski², Agnieszka Szymankiewicz-Szukała², Agnieszka Wincek²

¹**Oddział Neurologii, Szpital Wojewódzki w Koszalinie**

²**Zakład Patofizjologii Narządu Ruchu, Uniwersytet Medyczny w Poznaniu**

STRESZCZENIE

Spośród metod klinicznych oceny funkcji ruchowej u chorych po udarach niedokrwiennych mózgu, w świetle dostępnej literatury, niewiele umożliwia odpowiednie, zindywidualizowane wyznaczenie algorytmu usprawniania z wykorzystaniem metod kinezyterapii i fizykoterapii. Biorąc pod uwagę różnorodność obecnie stosowanych metod terapii, celem tego przeglądu jest przedstawienie nowoczesnych formy leczenia, opartych na przesłankach neurofizjologicznych, umożliwiających szybką oraz trwałą poprawę sprawności ruchowej u chorych po udarach, spełniając również funkcję narzędzia diagnostycznego, rzetelnie weryfikującego skuteczność terapii.

Słowa kluczowe: udar niedokrwienny, usprawnianie, diagnostyka, metody neurofizjologiczne

Data otrzymania: 19 luty 2019 Data zaakceptowania: 1 marzec 2019

Autorzy nie zgłosili źródła finansowania.

Autorzy nie deklarowali konfliktu interesów.

Autor odpowiedzialny za korespondencję:

Katarzyna Kaczmarek

Oddział Neurologii, Szpital Wojewódzki w Koszalinie

Chałubińskiego 7

75-950 Koszalin

gabinet@katarzynakaczmarek.pl

CURRENT NEUROPHYSIOLOGICAL METHODS FOR THE ASSESSMENT OF MOTOR FUNCTION AND THERAPY IN PATIENTS TREATED FOR ISCHEMIC STROKE

Katarzyna Kaczmarek¹, Piotr Brylewski², Agnieszka Szymankiewicz-Szukała², Agnieszka Wincek²

¹**Neurology Ward, Pomeranian District Hospital, Koszalin, Poland**

²**Department of Pathophysiology of Locomotor Organs, University of Medical Sciences, Poznań, Poland**

SUMMARY

Among the clinical methods of the motor function assessment in patients after ischemic stroke, in the light of the available literature, there is little to allow an appropriate, individualized determination of the rehabilitation algorithm using kinesitherapy and physical therapy

methods. Considering the diversity of currently used methods of therapy, the aim of this review is to present modern forms of treatment, based on neurophysiological aspects, enabling fast and permanent improvement of mobility in stroke patients, also fulfilling the function of a diagnostic tool that reliably verifies the effectiveness of therapy.

Keywords: ischemic stroke, rehabilitation, diagnostics, neurophysiological methods

Date received: 19th February 2019

Date accepted: 01st March 2019

Authors reported no source of funding.

Authors declared no conflict of interest.

Author responsible for correspondence (example):

Katarzyna Kaczmarek

Neurology Ward

Pomeranian District Hospital

Chałubińskiego 7

75-950 Koszalin

Poland

gabinet@katarzynakaczmarek.pl

Introduction

Stroke according to the World Health Organization (WHO) is a state of rapid, progressive, focal or generalized neurological deficits, life-threatening, leading to disability (Erajfej et al., 2017). Stroke may be caused by ischemia or bleeding in the central nervous system, and about 70-85% of strokes are ischemic. There are various qualifications of ischemic strokes due to the dynamics of symptoms, the area of ischemia and etiology (Giles and Rothwell, 2007).

Ischemic stroke can cause speech disorders, balance, cognitive, sensory and motor dysfunctions (Adams et al., 2003), depending on location of the ischemic area occupied, in the vicinity of the motor cortex or subcortical centers. In the practice of clinical post-stroke rehabilitation, a number of scales measuring the degree of disability and symptoms of loss of sensory and motor function are used, such as Hospital Anxiety and Depression Scale (HADS), Mini-Mental State Examination (MMSE), Modified Ashworth Scale (MAS), National Institutes of Health Stroke Scale (NIHSS), Stroke Rehabilitation Assessment of Movement (STREAM), Barthel Index (BI), Berg Balance Scale (BBS), Modified Rankin Handicap Scale (MRS), Motor Assessment Scale (MAS), 10-Meter Walking Test (Peural et al., 2002, Laufer et al., 2011, Sabut et al., 2011, Sahin et al., 2012, Erajfer et al., 2017, Guo et al., 2018). The results of evaluation of patients after ischemic stroke using the assessment scales listed above, however, do not provide indications for the introduction of specific therapeutic procedures for kinesitherapy and physical therapy as well as pharmacological treatment (ESO Guidelines, 2008).

Aim

Considering the diversity of currently used methods of therapy, the aim of this review was to present modern forms of treatment based on neurophysiological aspects, enabling fast and permanent improvement of mobility in stroke patients, also fulfilling the function of a diagnostic tool that reliably verifies the effectiveness of therapy.

Methods

Pharmacologically, most patients with well-recognised ischemic stroke are treated with antiplatelet drugs, mainly acetylsalicylic acid, and in cases of contraindications to treatment with ASA, thienopyridine drugs such as clopidogrel are used. In a patient admitted to hospital

treatment in a so-called "therapeutic window" (that is, up to 4.5 hours from the onset of symptoms) and without contraindications to intravenous injection of tissue plasminogen activator (Alteplase). In special cases, mechanical thrombectomy may be used during acute ischemic stroke due to large vessel intracranial occlusion.


Modern physiotherapeutic treatment of patients after ischemic stroke is based on the assumption of relearning the motor functions related to neuroplasticity, both at the level of spinal cord centers activity and at the supraspinal level centers (Woldag and Hummelsheim 2002, Wolf et al., 2005). Therefore, the most commonly used form of kinesitherapy is PNF treatment, also called proprioceptive neuromuscular facilitation (Woldag and Hummelsheim 2002). The other treatments are individualized (Auchstaetter et al., 2016), depending on the patient's psychophysical condition, but mainly include passive, supportive and active extremities exercises on the side of the paresis, exercises reducing the spasticity symptom (PIR - post-isometric relaxation procedures), stretching exercises, as well as procedures stimulating proprioceptors. During the 10-day stay in the post-stroke rehabilitation ward, the patient is up righted to standing position with also training to sit down, teaching changes in position and global movements that are performed during everyday life (e.g. changing positions in bed, moving from a chair to a chair). Another element of rehabilitation is learning to walk (locomotion), including gait at the handrails and walk using orthopedic equipment ('balcony' support). Other exercises are respiratory and equivalent, in which the eyesight, hearing, receptors of touch, balance and deep sensation are activated.

Furthermore, physiotherapeutic treatment mainly include "warming therapy" and electrostimulation of paralysed muscles using the algorithms of electrical stimuli normally supplied with electrotherapy equipment by manufacturers (Hara 2013, Stein et al., 2015). The adaptation of the latter leaves much to be desired when it comes to individualization for the needs of the rehabilitated patient. Electrotherapy is sometimes enriched with functional stimulation of the muscles acting antagonistically at the level of the wrist and ankle joint (Lisiński et al., 2008). In the studies of Lisiński et al. (2008), similarly to the studies of Kraft et al. (1992), who used combination of PNF and functional electrical stimulation (FES) in stroke patients, a significant improvement in the function of weak muscles of the symptomatic extremities was found, both upper and lower. Until now, however, studies in this area were performed in relation to a small group of patients after ischemic stroke (N = 24) in short-term observation (after 20 days of treatment) (Lisiński et al. 2008). In addition to electrostimulation treatments that are supposed to affect the efficiency of motor units or muscles themselves or improve the conductivity of nerve impulses in motor nerve fibers (Sharififar et al., 2018), neuroplastic phenomena have been documented at the level of the sensory cortex after stimulation of sensory nerve fibers (Peurala et al. , 2002, Laufer et al., 2011). Other physical therapy procedures include treatments in the field of thermotherapy and electrostimulation that should induce lowering of muscle tone in patients after stroke with a symptom of spasticity (Collins et al., 1986, Bell and Lermann 1987, Shields 2002, Garcia et al., 2019).

One of the newest methods of therapy of patients after ischemic stroke is using the repetitive transcranial magneticstimulation with a series of magnetic stimuli (rTMS). The research on this therapy was conducted by Guan et al (2017), who showed an improvement in motor function and a reduction of spasticity symptoms.

For a neurologist, rehabilitation doctor or physiotherapist, in a professional practice, a method perfectly suited to the treatment of ischemic stroke would be one that would allow precise quantification of muscle function and nerve impulses conduction, and on the other would be based on results obtained in a targeted manner from a patient's diagnostic tests. A proposition in the field of clinical neurophysiology, in conjunction with the principle of muscle electrostimulation based on the biofeedback principle, is the mentioned remarkable regulation of muscular actions acting antagonistically on the wrist joints and the ankle joints. The alternating muscular activity is regulated by the Ia inhibitory interneurone spinal cord system (Bhagchandani and Schindler-Ivens, 2012). In stroke patients, it is released from the control of

supernatural centers. The electromyograms recorded with surface electrodes (sEMG) show disturbances, the range of which can be indexed on a scale of 5-1 (5 - alternating normal activity, 1 - no alternation) (Figure 1).



Alternate activity of antagonistic muscles acting at the same joint




Figure 1. Recordings of electromyograms showing changes in motor unit function of antagonistic muscles of various degrees of severity (a- extensor carpii and flexores carpi muscles, b- tibialis anterior and gastrocnemius muscles) acting at the same joint in patients after stroke scored from 5 to 1 (A - (5) alternate, proper activity; E, F - (1) no alternation) (Lisiński et al 2008, with permission).

In patients after stroke, functional electrostimulation of the antagonistic muscles acting on the wrist and ankle joints can be used (Figure 2) using a personal, portable, 4-channel electrostimulation device.



Figure 2. Photo of the apparatus for electrostimulation of antagonistic muscles acting on the wrist joint with the presentation of the location of stimulating electrodes. Source: Department of Pathophysiology of Locomotor Organs, University Medical Sciences in Poznań.

The algorithm can be individually adapted for a patient with certain parameters based on previous sEMG and electroneurographic tests of nerve impulses in motor fibers, depending on the severity of pathological changes and the frequency of sEMG recordings during attempts of maximal contraction condition (usually at 40-70Hz). Contraindications to therapy are epileptic episodes and other consequences of direct craniocerebral injuries, serious cardiovascular dysfunction, pregnancy, ferromagnetic implants, electronic implants such as a pacemaker or a cochlear implant.

Conclusions

The modern methods of rehabilitation of stroke patients applied so far, in the majority of cases, are based on neurophysiological records, including those that use biofeedback, which seem to be the most suitable, because they meet the criteria of individualization in relation to the rehabilitated patient. Future studies verifying the effectiveness of electrostimulation procedures should focus on combining them with targeted kinesitherapy (Adams et al., 2003; Duncac et al., 2005).

REFERENCES

- Adams H.P. Jr, Adams R.J., Brott T., del Zoppo G.J., Furlan A., Goldstein L.B.** (2003) „*Guidelines for the early management of patients with ischemic stroke: a scientific statement from the Stroke Council of the American Stroke Association.*” *Stroke*. 34, pp. 1056–1083.
- Auchstaetter N, Luc J., Lukye S., Lynd K., Schemenauer S., Whittaker M., Musselman K.E.** (2016) “*Use of Functional Electrical Stimulation for Clients With Stroke: Frequency, Barriers, and Facilitators.*” *Physical Therapy*, 96(7), pp. 995–1005,
- Bell K.R., Lermann J.F.** (1987) “*Effect of cooling on H and T - reflexes in normal subjects.*” *Arch Psych Med Rehabil*, 45(68), pp. 490-501.
- Bhagchandani N., Schindler-Ivens S.** (2012) “*Reciprocal inhibition post-stroke is related to reflex excitability and movement ability.*” *Clin Neurophysiol*. 123(11), pp. 2239–2246.
- Kraft G.H., Fitas S.S., Hammond M.C.** (1992) “*Techniques to improve function of the arm and hand in chronic hemiplegia.*” *Arch Phys Med Rehabil*, 73, pp. 220-227.
- Collins K., Story M., Peterson K.** (1986) “*Peroneal nerves palsy after cryotherapy.*” *Phys Sports Med*, 14, pp. 105-108
- Duncan P.W., Zorowitz R., Bates B.** (2005) “*Management of Adult Stroke Rehabilitation Care: a clinical practice guideline.*” *Stroke*., 36, pp. 100–143.
- Eraifej J., Clark W., France B, Desando S., Moore D.** (2017) “*Effectiveness of upper limb functional electrical stimulation after stroke for the improvement of activities of daily living and motor function: a systematic review and meta-analysis.*” *Systematic Reviews* 6:40
- Garcia L.C., Alcântara C.C., Santos G.L., Monção J.V.A., Russo T.L.** (2019) “*Cryotherapy Reduces Muscle Spasticity But Does Not Affect Proprioception in Ischemic Stroke: A Randomized Sham-Controlled Crossover Study.*” *Am J Phys Med Rehabil*, 98(1), pp. 51-57.
- Giles M.F., Rothwell P.M.** (2007) “*Risk of stroke early after transient ischaemic attack: a systematic review and meta-analysis.*” *Lancet Neurol*, 6, pp. 1063-1107.
- Guan Y.Z., Li J., Zhang X.W., Wu S., Du H., Cui L.Y., Zhang W.H.** (2017) “*Effectiveness of repetitive transcranial magnetic stimulation (rTMS) after acute stroke: A one-year longitudinal randomized trial.*” *CNS Neurosci Ther.*, 23(12), pp. 940-946.
- Guo X-x., Fan B-y., Mao Y-y.** (2018) “*Effectiveness of neuromuscular electrical stimulation for wrist rehabilitation after acute ischemic stroke.*” *Medicine*. 97(38), pp. e12299.
- Hara Y.** (2013) “**Rehabilitation with Functional Electrical Stimulation in Stroke Patients**” *Int J Phys Med Rehabil*, 1(6)
- Laufer Y., Elboim-Gabyzon M.** “*Does Sensory Transcutaneous Electrical Stimulation Enhance Motor Recovery Following a Stroke?*” A Systematic Review. *Neurorehabilitation and Neural Repair*, 25(9), pp. 799 –809.

- Lisiński P., Huber J., Samborski W., Witkowska A.** (2008) *“Neurophysiological assessment of the electrostimulation procedures used in stroke patients during rehabilitation.”* Int. J. Artif. Organs., 31(1), pp. 76-86.
- Peurala S.H., Pitkänen K., Sivenius J., Tarkka I.M.** (2002) *„Cutaneous electrical stimulation may enhance sensorimotor recovery in chronic stroke”* Clinical Rehabilitation., 1(6), pp. 709–716.
- Sabut S.K., Sikdar C., Kumar R., Mahadevappa M.** (2011) *“Functional electrical stimulation of dorsiflexor muscle: effects on dorsiflexor strength, plantarflexor spasticity, and motor recovery in stroke patients”*. NeuroRehabilitation., 29(4), pp. 393-400.
- Sahin N., Ugurlu H., Albayrak I.** (2012) *“The efficacy of electrical stimulation in reducing the post-stroke spasticity: a randomized controlled study”* Disability and Rehabilitation, 34:2, pp. 151-156,
- Shields N.** (2002) *“Short -wave diathermy: current clinical and safety practices.”* Physiotherapy Research International, 7(4), pp. 191-202.
- Stein C., Fritsch C.G., Robinson C., Sbruzzi G., Della R., Plentz M.** (2015) *“Effects of Electrical Stimulation in Spastic Muscles After Stroke. Systematic Review and Meta-Analysis of Randomized Controlled Trials”*. Stroke., 46, pp. 2197-2205.
- The European Stroke Organisation (ESO) Executive Committee and the ESO Writing Committee Guidelines for Management of Ischaemic Stroke and Transient Ischaemic Attack* (2008). Cerebrovasc. 25: pp. 457–507.
- Woldag H., Hummelsheim H.** (2002) *“Evidence-based physiotherapeutic concepts for improving arm and hand function in stroke patients: a review.”* J Neurol., 249, pp. 518-528.
- Wolf S.L., Butler A.J., Alberts J.L., Kim M.W.** (2005) *“Contemporary linkages between EMG kinetics and stroke rehabilitation.”* Journal of electromyography and kinesiology., 15, pp. 229-239.